A speedboat with a brown canvas top is moving across the water, leaving a white wake. In the background, a dense city skyline with various skyscrapers is visible under a hazy sky. A line of green trees separates the water from the city.

Dyson-Schwinger Equations and Quantum Chromodynamics

Craig D. Roberts
`cdroberts@anl.gov`

Physics Division
Argonne National Laboratory

<http://www.phy.anl.gov/theory/staff/cdr.html>

Craig Roberts, Dyson-Schwinger Equations and QCD
in Physics: Workshop on Electromagnetic Interactions, 34/60 - US DOE, 2006... - p. 1/38

cdroberts@anl.gov

Argonne National Laboratory


<http://www.phy.anl.gov/theory/staff/cdr.html>

Room with a View




Universal Truths

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Universal Truths


[First](#)[Contents](#)[Back](#)[Conclusion](#)



Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.






Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach.






Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach. Covariance requires existence of quark orbital angular momentum in hadron's rest-frame wave function.






Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach. Covariance requires existence of quark orbital angular momentum in hadron's rest-frame wave function.
- DCSB is most important mass generating mechanism for matter in the Universe.






Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach. Covariance requires existence of quark orbital angular momentum in hadron's rest-frame wave function.
- DCSB is most important mass generating mechanism for matter in the Universe. Higgs mechanism is irrelevant to light-quarks.





Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach. Covariance requires existence of quark orbital angular momentum in hadron's rest-frame wave function.
- DCSB is most important mass generating mechanism for matter in the Universe. Higgs mechanism is irrelevant to light-quarks.
- Challenge: understand relationship between parton properties on the light-front and rest frame structure of hadrons.





Universal Truths

- Form factors give information about distribution of hadron's characterising properties amongst its QCD constituents.
- Calculations at $Q^2 > 1 \text{ GeV}^2$ require a Poincaré-covariant approach. Covariance requires existence of quark orbital angular momentum in hadron's rest-frame wave function.
- DCSB is most important mass generating mechanism for matter in the Universe. Higgs mechanism is irrelevant to light-quarks.
- Challenge: understand relationship between parton properties on the light-front and rest frame structure of hadrons. Problem because, e.g., DCSB - an established keystone of low-energy QCD and the origin of constituent-quark masses - has not been realised in the light-front formulation.



Form Factors: Why?

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Form Factors: Why?

- The nucleon and pion hold special places in non-perturbative studies of QCD.



Form Factors: Why?

- The nucleon and pion hold special places in non-perturbative studies of QCD.
- An explanation of nucleon and pion structure and interactions is central to hadron physics – they are respectively the archetypes for baryons and mesons.



Form Factors: Why?

- The nucleon and pion hold special places in non-perturbative studies of QCD.
- An explanation of nucleon and pion structure and interactions is central to hadron physics – they are respectively the archetypes for baryons and mesons.
- Form factors have long been recognized as a basic tool for elucidating bound state properties. They can be studied from very low momentum transfer, the region of non-perturbative QCD, up to a region where perturbative QCD predictions can be tested.



Form Factors: Why?

- The nucleon and pion hold special places in non-perturbative studies of QCD.
- An explanation of nucleon and pion structure and interactions is central to hadron physics – they are respectively the archetypes for baryons and mesons.
- Form factors have long been recognized as a basic tool for elucidating bound state properties. They can be studied from very low momentum transfer, the region of non-perturbative QCD, up to a region where perturbative QCD predictions can be tested.
- Experimental and theoretical studies of nucleon electromagnetic form factors have made rapid and significant progress during the last several years, including new data in the time like region, and material gains have been made in studying the pion form factor.



Form Factors: Why?

- The nucleon and pion hold special places in non-perturbative studies of QCD.
- An explanation of nucleon and pion structure and interactions is central to hadron physics – they are respectively the archetypes for baryons and mesons.
- Form factors have long been recognized as a basic tool for elucidating bound state properties. They can be studied from very low momentum transfer, the region of non-perturbative QCD, up to a region where perturbative QCD predictions can be tested.
- Experimental and theoretical studies of nucleon electromagnetic form factors have made rapid and significant progress during the last several years, including new data in the time like region, and material gains have been made in studying the pion form factor.
- Despite this, many urgent questions remain unanswered.



Some Questions

- What is the role of pion cloud in nucleon electromagnetic structure?
- Can we understand the pion cloud in a more quantitative and, perhaps, model-independent way?



Some Questions

- Where is the transition from non-pQCD to pQCD in the pion and nucleon electromagnetic form factors?



Some Questions

- Do we understand the high Q^2 behavior of the proton form factor ratio in the space-like region?
- Can we make model-independent statements about the role of relativity or orbital angular momentum in the nucleon?



Some Questions

- Can we understand the rich structure of the time-like proton form factors in terms of resonances?
- What do we expect for the proton form factor ratio in the time-like region?
- What is the relation between proton and neutron form factor in the time-like region?
- How do we understand the ratio between time-like and space-like form factors?



Some Questions

- What is the role of two-photon exchange contributions in understanding the discrepancy between the polarization and Rosenbluth measurements of the proton form factor ratio?
- What is the impact of these contributions on other form factor measurements?



Some Questions

- How accurately can the pion form factor be extracted from the $ep \rightarrow e'n\pi^+$ reaction?





- Current status is described in
 - J. Arrington, C. D. Roberts and J. M. Zanotti
“Nucleon electromagnetic form factors,”
J. Phys. G **34**, S23 (2007); [arXiv:nucl-th/0611050].
 - C. F. Perdrisat, V. Punjabi and M. Vanderhaeghen,
“Nucleon electromagnetic form factors,”
Prog. Part. Nucl. Phys. **59**, 694 (2007);
[arXiv:hep-ph/0612014].




- Current status is described in
 - J. Arrington, C. D. Roberts and J. M. Zanotti
“Nucleon electromagnetic form factors,”
J. Phys. G **34**, S23 (2007); [arXiv:nucl-th/0611050].
 - C. F. Perdrisat, V. Punjabi and M. Vanderhaeghen,
“Nucleon electromagnetic form factors,”
Prog. Part. Nucl. Phys. **59**, 694 (2007);
[arXiv:hep-ph/0612014].





- Most recently:
“ECT* Workshop on Hadron Electromagnetic Form Factors”
Organisers: Alexandrou, Arrington, Friedrich, Maas, Roberts
Presentations, etc., available on-line
<http://ect08.phy.anl.gov/>

Answer for the pion




Answer for the pion

Two \rightarrow Infinitely many ...



Answer for the pion

Two \rightarrow Infinitely many ...
Handle that
properly in
quantum
field theory




Answer for the pion

Two \rightarrow Infinitely many ...

Handle that properly in quantum field theory

...

momentum-dependent dressing



Answer for the pion

Two \rightarrow Infinitely many ...


Handle that properly in quantum field theory

...

momentum-dependent dressing

...

perceived distribution of mass depends on the resolving scale



Pion Form Factor

Procedure Now Straightforward



[First](#)


[Contents](#)

[Back](#)

[Conclusion](#)


Pion Form Factor

- Solve Gap Equation
⇒ Dressed-Quark Propagator, $S(p)$




Pion Form Factor

- Use that to Complete Bethe Salpeter Kernel, K
- Solve Homogeneous Bethe-Salpeter Equation for Pion Bethe-Salpeter Amplitude, Γ_π



Pion Form Factor

- Use that to Complete Bethe Salpeter Kernel, K
- Solve Homogeneous Bethe-Salpeter Equation for Pion Bethe-Salpeter Amplitude, Γ_π




- Solve Inhomogeneous Bethe-Salpeter Equation for Dressed-Quark-Photon Vertex, Γ_μ




Pion Form Factor

- Now have all elements for Impulse Approximation to Electromagnetic Pion Form factor



Pion Form Factor

- Now have all elements for Impulse Approximation to Electromagnetic Pion Form factor




- Evaluate this final, three-dimensional integral



Calculated Pion Form Factor

Calculation first published in 1999; No Parameters Varied

Numerical method improved in 2005




Calculated Pion Form Factor

Calculation first published in 1999; No Parameters Varied

Numerical method improved in 2005

Data published
in 2001.

Subsequently
revised





Timelike Pion Form Factor

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Timelike Pion Form Factor

Ab initio calculation into timelike region

Deeper than ground-state ρ -meson pole






Timelike Pion Form Factor

Ab initio calculation into timelike region

Deeper than ground-state ρ -meson pole






Timelike Pion Form Factor

Ab initio calculation into timelike region

Deeper than ground-state ρ -meson pole

ρ -meson not put in “by hand” – generated dynamically as a bound-state of dressed-quark and dressed-antiquark



Dimensionless product: $r_\pi f_\pi$





Dimensionless product: $r_\pi f_\pi$

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction





Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation





Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Great strides towards placing nucleon studies on same footing as mesons



Dimensionless product: $r_\pi f_\pi$


- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Experimentally: $r_\pi f_\pi = 0.315 \pm 0.005$



Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Experimentally: $r_\pi f_\pi = 0.315 \pm 0.005$

● DSE prediction




Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Experimentally: $r_\pi f_\pi = 0.315 \pm 0.005$

DSE prediction

Lattice results


– James Zanotti [UK QCD]



Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Experimentally: $r_\pi f_\pi = 0.315 \pm 0.005$


- DSE prediction
- Lattice results
 - James Zanotti [UK QCD]
- Fascinating result:
DSE and Lattice
 - Experimental value obtains independent of current-quark mass.



Dimensionless product: $r_\pi f_\pi$

- Improved rainbow-ladder interaction
- Repeating $F_\pi(Q^2)$ calculation
- Experimentally: $r_\pi f_\pi = 0.315 \pm 0.005$

- DSE prediction
- Fascinating result:
DSE and Lattice
– Experimental value
obtains independent of
current-quark mass.
We have understood this
Implications far-reaching.





New Challenges

[First](#)[Contents](#)[Back](#)[Conclusion](#)

New Challenges

- **Next Steps** . . . Applications to excited states and axial-vector mesons, e.g., will improve understanding of confinement interaction between light-quarks.




New Challenges

- **Next Steps** . . . Applications to excited states and axial-vector mesons, e.g., will improve understanding of confinement interaction between light-quarks.
- Move on to the problem of a **symmetry preserving** treatment of hybrids and exotics.



New Challenges


- Another Direction . . . Also want/need information about three-quark systems



New Challenges

- Another Direction . . . Also want/need information about three-quark systems

- With this problem . . . most wide-ranging studies employ expertise familiar from meson applications circa ~ 1995 .




New Challenges

- Another Direction . . . Also want/need information about three-quark systems


- With this problem . . . most wide-ranging studies employ expertise familiar from meson applications circa ~ 1995 .

- Namely . . . Model-building and Phenomenology, constrained by the DSE results outlined already.



New Challenges

- Another Direction . . . Also want/need information about three-quark systems




- With this problem . . . most wide-ranging studies employ expertise familiar from meson applications circa ~ 1995 .
- However, that is beginning to change . . .



New Challenges


- Another Direction . . . Also want/need information about three-quark systems



- With this problem . . . most wide-ranging studies employ expertise familiar from meson applications circa ~ 1995 .
- However, that is beginning to change . . .




Nucleon ... Three-body Problem?



Nucleon ...

Three-body Problem?


- What is the picture in quantum field theory?



Nucleon ...

Three-body Problem?


- What is the picture in quantum field theory?
- Three → infinitely many!





Faddeev equation

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Faddeev equation




Faddeev equation



- Linear, Homogeneous Matrix equation
 - Yields *wave function* (Poincaré Covariant Faddeev Amplitude) that describes quark-diquark relative motion within the nucleon
- Scalar and Axial-Vector Diquarks ... In Nucleon's Rest Frame Amplitude has ... s -, p - & d -wave correlations

Diquark correlations



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)

QUARK-QUARK

Craig Roberts: Dyson Schwinger Equations and QCD

25th Students' Workshop on Electromagnetic Interactions, 31/08 – 05/09, 2008. ... – p. 16/38

Diquark correlations

- Same interaction that describes mesons also generates three coloured quark-quark correlations: blue–red, blue–green, green–red


- Confined ... Does not escape from within baryon.


- Scalar is isosinglet, Axial-vector is isotriplet

- DSE and lattice-QCD

$$m_{[ud]_{0+}} = 0.74 - 0.82$$

$$m_{(uu)_{1+}} = m_{(ud)_{1+}} = m_{(dd)_{1+}} = 0.95 - 1.02$$





Nucleon EM Form Factors: A Précis

Höll, et al.: nu-th/0412046 & nu-th/0501033




[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)



Nucleon EM Form Factors: A Précis

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Nucleon EM Form Factors: A Précis

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)

Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)

Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)

Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

(Oettel, Hellstern, Alkofer, Reinhardt: [nucl-th/9805054](#))



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)

Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

- But is that good?



Nucleon EM Form Factors: A Précis

Cloët, et al.:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)

Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

- But is that good?
 - Cloudy Bag: $\delta M_+^{\pi\text{-loop}} = -300$ to -400 MeV!



Nucleon EM Form Factors: A Précis

Cloët, *et al.*:

arXiv:0710.2059, arXiv:0710.5746 & arXiv:0804.3118

- Interpreting expts. with GeV electromagnetic probes requires Poincaré covariant treatment of baryons
⇒ Covariant dressed-quark Faddeev Equation
- Excellent mass spectrum (octet and decuplet)

Easily obtained:

$$\left(\frac{1}{N_H} \sum_H \frac{[M_H^{\text{exp}} - M_H^{\text{calc}}]^2}{[M_H^{\text{exp}}]^2} \right)^{1/2} = 2\%$$

- But is that good?
 - Cloudy Bag: $\delta M_+^{\pi\text{-loop}} = -300$ to -400 MeV!
- Critical to anticipate pion cloud effects

Roberts, Tandy, Thomas, *et al.*, nu-th/02010084



Nucleon's self-energy - pion loop



Nucleon's self-energy - pion loop

$$\begin{aligned}\Sigma(P) &= 3 \int \frac{d^4 k}{(2\pi)^4} g_{PV}^2(P, k) \Delta_\pi((P - k)^2) \\ &\quad \times \boxed{\gamma \cdot (P - k) \gamma_5} G(k) \boxed{\gamma \cdot (P - k) \gamma_5} \\ &= i\gamma \cdot k [\mathcal{A}(k^2) - 1] + \mathcal{B}(k^2)\end{aligned}$$



Nucleon's self-energy - pion loop

$$\begin{aligned}\Sigma(P) &= 3 \int \frac{d^4 k}{(2\pi)^4} g_{PV}^2(P, k) \Delta_\pi((P - k)^2) \\ &\quad \times \boxed{\gamma \cdot (P - k) \gamma_5} G(k) \boxed{\gamma \cdot (P - k) \gamma_5} \\ &= i\gamma \cdot k [\mathcal{A}(k^2) - 1] + \mathcal{B}(k^2)\end{aligned}$$

- Pseudovector coupling



Nucleon's self-energy - pion loop

$$\begin{aligned}\Sigma(P) &= 3 \int \frac{d^4 k}{(2\pi)^4} g_{PV}^2(P, k) \Delta_\pi((P - k)^2) \\ &\quad \times \boxed{\gamma \cdot (P - k) \gamma_5} G(k) \boxed{\gamma \cdot (P - k) \gamma_5} \\ &= i\gamma \cdot k [\mathcal{A}(k^2) - 1] + \mathcal{B}(k^2)\end{aligned}$$

- Pseudovector coupling
- Completely equivalent to pseudoscalar coupling IF that is treated completely
- Tadpole contribution **can't** be neglected

(Hecht, Oettel, Roberts, Schmidt, Tandy, Thomas: [nucl-th/0201084](#))



Nucleon's self-energy - pion loop

$$\begin{aligned}\Sigma(P) &= 3 \int \frac{d^4 k}{(2\pi)^4} g_{PV}^2(P, k) \Delta_\pi((P - k)^2) \\ &\quad \times \boxed{\gamma \cdot (P - k) \gamma_5} G(k) \boxed{\gamma \cdot (P - k) \gamma_5} \\ &= i\gamma \cdot k [\mathcal{A}(k^2) - 1] + \mathcal{B}(k^2)\end{aligned}$$



$g_{PV}(P, k)$, πN vertex function



Calculated using Γ_π and Ψ_N



Always soft: Monopole $\lambda \sim 0.6$ GeV



Nucleon's self-energy - pion loop

$$\begin{aligned}\Sigma(P) &= 3 \int \frac{d^4 k}{(2\pi)^4} g_{PV}^2(P, k) \Delta_\pi((P-k)^2) \\ &\quad \times \boxed{\gamma \cdot (P-k) \gamma_5} G(k) \boxed{\gamma \cdot (P-k) \gamma_5} \\ &= i\gamma \cdot k [\mathcal{A}(k^2) - 1] + \mathcal{B}(k^2)\end{aligned}$$



$g_{PV}(P, k)$, πN vertex function



Calculated using Γ_π and Ψ_N




Always soft: Monopole $\lambda \sim 0.6$ GeV



Corresponds to range $r_\lambda \sim 0.8$ fm

... pion cloud does not

penetrate deeply within nucleon.



Nucleon's self-energy - pion loop

$$\begin{aligned}\Sigma(P) &= 3 \int \frac{d^4 k}{(2\pi)^4} g_{PV}^2(P, k) \Delta_\pi((P - k)^2) \\ &\quad \times \boxed{\gamma \cdot (P - k) \gamma_5} G(k) \boxed{\gamma \cdot (P - k) \gamma_5} \\ &= i\gamma \cdot k [\mathcal{A}(k^2) - 1] + \mathcal{B}(k^2)\end{aligned}$$



$$\begin{aligned}G(k) &= 1/[i\gamma \cdot k + M + \Sigma(P)] && \text{Pole Position Not} \\ &= -i\gamma \cdot k \sigma_V(k^2) + \sigma_S(k^2) && \text{Known a priori}\end{aligned}$$

● Mass shift calculated via self-consistent solution

Nucleon's self-energy - pion loop

$$\begin{aligned}\Sigma(P) &= 3 \int \frac{d^4 k}{(2\pi)^4} g_{PV}^2 (\text{const.}) \Delta_\pi((P-k)^2) \\ &\quad \times \boxed{\gamma \cdot (P-k) \gamma_5} G(k) \boxed{\gamma \cdot (P-k) \gamma_5} \\ &= i\gamma \cdot k [\mathcal{A}(k^2) - 1] + \mathcal{B}(k^2)\end{aligned}$$



Argonne
NATIONAL
LABORATORY

Obtain Integral Equation Kernels

$$\int d\Omega_k f((P-k)^2) = \frac{2}{\pi} \int_{-1}^1 dz \sqrt{1-z^2} f(P^2 + k^2 - 2P kz)$$

E.g.

$$\omega_B(P^2, k^2) = \int d\Omega_k \frac{(P-k)^2}{(P-k)^2 + m_\pi^2} = 1 - \frac{2m_\pi^2}{a + \sqrt{a^2 - b^2}},$$

$$a = P^2 + k^2 + m_\pi^2, b = 2Pk$$

Craig Roberts: Dyson-Schwinger Equations and QCD

25th Students' Workshop on Electromagnetic Interactions, 31/08 – 05/09, 2008. ... – p. 19/38

First

Contents

Back

Conclusion

Nucleon's self-energy - pion loop

$$\begin{aligned}\Sigma(P) &= 3 \int \frac{d^4 k}{(2\pi)^4} g_{PV}^2(P, k) \Delta_\pi((P - k)^2) \\ &\quad \times \boxed{\gamma \cdot (P - k) \gamma_5} G(k) \boxed{\gamma \cdot (P - k) \gamma_5} \\ &= i\gamma \cdot k [\mathcal{A}(k^2) - 1] + \mathcal{B}(k^2)\end{aligned}$$



● But $g_{PV} = g_{PV}(P^2, k^2, P \cdot k)$

Therefore, **In General**, Kernel only known Numerically

● Complicates analysis ...

locating, incorporating poles in integrand

Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)

Nucleon Self Energy: Chiral Limit

Hecht, *et al.*, nu-th/0201084

- Let's look what happens when $m_\pi \rightarrow 0$
 - Minkowski Space
 - Pseudovector Coupling



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

- Let's look what happens when $m_\pi \rightarrow 0$
 - Minkowski Space
 - Pseudovector Coupling
- One-loop nucleon self energy

$$\Sigma(P) = 3i \frac{g^2}{4M^2} \int \frac{d^4 k}{(2\pi)^4} \Delta(k^2, m_\pi^2) \not{k} \gamma_5 G_0(P - k) \not{k} \gamma_5 .$$

This integral is divergent. Assume a Poincaré covariant regularisation, characterised by a mass-scale λ



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

- Let's look what happens when $m_\pi \rightarrow 0$
 - Minkowski Space
 - Pseudovector Coupling
- One-loop nucleon self energy

$$\Sigma(P) = 3i \frac{g^2}{4M^2} \int \frac{d^4 k}{(2\pi)^4} \Delta(k^2, m_\pi^2) \not{k} \gamma_5 G_0(P - k) \not{k} \gamma_5 .$$

This integral is divergent. Assume a Poincaré covariant regularisation, characterised by a mass-scale λ

- Decompose nucleon propagator into positive and negative energy components

$$\begin{aligned} G_0(P) &= \frac{1}{\not{P} - M_0} = G_0^+(P) + G_0^-(P) \\ &= \frac{M}{\omega_N(\vec{P})} \left[\Lambda_+(\vec{P}) \frac{1}{P_0 - \omega_N(\vec{P}) + i\varepsilon} + \Lambda_-(\vec{P}) \frac{1}{P_0 + \omega_N(\vec{P}) - i\varepsilon} \right] \quad (4) \end{aligned}$$

$$\omega_N^2(\vec{P}) = \vec{P}^2 + M^2, \text{ and } \Lambda_\pm(\vec{P}) = (\tilde{P} \pm M)/(2M), \tilde{P} = (\omega(\vec{P}), \vec{P})$$



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)

Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

- One-loop nucleon self energy

$$\Sigma(P) = 3i \frac{g^2}{4M^2} \int \frac{d^4 k}{(2\pi)^4} \Delta(k^2, m_\pi^2) \not{k} \gamma_5 G_0(P - k) \not{k} \gamma_5 .$$



Nucleon Self Energy: Chiral Limit

Hecht, *et al.*, nu-th/0201084

- One-loop nucleon self energy

$$\Sigma(P) = 3i \frac{g^2}{4M^2} \int \frac{d^4 k}{(2\pi)^4} \Delta(k^2, m_\pi^2) \not{k} \gamma_5 G_0(P - k) \not{k} \gamma_5 .$$

- Shift in the mass of a positive energy nucleon nucleon:
$$\delta M_+ = \frac{1}{2} \text{tr}_D \left[\Lambda_+(\vec{P} = 0) \Sigma(P_0 = M, \vec{P} = 0) \right]$$



Nucleon Self Energy: Chiral Limit

Hecht, *et al.*, nu-th/0201084

- One-loop nucleon self energy

$$\Sigma(P) = 3i \frac{g^2}{4M^2} \int \frac{d^4 k}{(2\pi)^4} \Delta(k^2, m_\pi^2) \not{k} \gamma_5 G_0(P - k) \not{k} \gamma_5 .$$

- Shift in the mass of a positive energy nucleon nucleon:
$$\delta M_+ = \frac{1}{2} \text{tr}_D \left[\Lambda_+(\vec{P} = 0) \Sigma(P_0 = M, \vec{P} = 0) \right]$$
- Focus on positive energy nucleon's contribution to the loop integral; i.e., $\Delta(k) G^+(P - k)$, which we denote: $\delta_F M_+^+$



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

- One-loop nucleon self energy

$$\Sigma(P) = 3i \frac{g^2}{4M^2} \int \frac{d^4 k}{(2\pi)^4} \Delta(k^2, m_\pi^2) \not{k} \gamma_5 G_0(P - k) \not{k} \gamma_5 .$$

- Shift in the mass of a positive energy nucleon nucleon:

$$\delta M_+ = \frac{1}{2} \text{tr}_D \left[\Lambda_+(\vec{P} = 0) \Sigma(P_0 = M, \vec{P} = 0) \right]$$

- Focus on positive energy nucleon's contribution to the loop integral; i.e., $\Delta(k) G^+(P - k)$, which we denote: $\delta_F M_+^+$

- To evaluate k_0 integral, close contour in lower half-plane, thereby encircling only the positive-energy pion pole.

$$\delta_F M_+^+ = -3g^2 \int \frac{d^3 k}{(2\pi)^3} \frac{\omega_N(\vec{k}^2) - M_0}{4 \omega_N(\vec{k}^2)} \frac{1}{\omega_\pi(\vec{k}^2) [\omega_\pi(\vec{k}^2) + \omega_N(\vec{k}^2) - M_0]} \quad (9)$$



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

$$\delta_F M_+^+ = -3g^2 \int \frac{d^3 k}{(2\pi)^3} \frac{\omega_N(\vec{k}^2) - M_0}{4 \omega_N(\vec{k}^2)} \frac{1}{\omega_\pi(\vec{k}^2) [\omega_\pi(\vec{k}^2) + \omega_N(\vec{k}^2) - M_0]} \quad (10)$$



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

$$\delta_F M_+^+ = -3g^2 \int \frac{d^3 k}{(2\pi)^3} \frac{\omega_N(\vec{k}^2) - M_0}{4 \omega_N(\vec{k}^2)} \frac{1}{\omega_\pi(\vec{k}^2) [\omega_\pi(\vec{k}^2) + \omega_N(\vec{k}^2) - M_0]} \quad (14)$$

- On the domain for which the regularised integral has significant support, assume that M_0 is very much greater than all other mass scales.

$$\omega_N(\vec{k}^2) - M \approx \frac{\vec{k}^2}{2M} \quad (15)$$



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

$$\delta_F M_+^+ = -3g^2 \int \frac{d^3 k}{(2\pi)^3} \frac{\omega_N(\vec{k}^2) - M_0}{4 \omega_N(\vec{k}^2)} \frac{1}{\omega_\pi(\vec{k}^2) [\omega_\pi(\vec{k}^2) + \omega_N(\vec{k}^2) - M_0]} \quad (18)$$

- On the domain for which the regularised integral has significant support, assume that M_0 is very much greater than all other mass scales.

$$\omega_N(\vec{k}^2) - M \approx \frac{\vec{k}^2}{2M} \quad (19)$$

- Then

$$\delta_F M_+^+ \approx -3g^2 \int \frac{d^3 k}{(2\pi)^3} \frac{\vec{k}^2}{8M^2 \omega_{\lambda_i}^2(\vec{k}^2)} \quad (20)$$



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

$$\delta_F M_+^+ = -3g^2 \int \frac{d^3 k}{(2\pi)^3} \frac{\omega_N(\vec{k}^2) - M_0}{4 \omega_N(\vec{k}^2)} \frac{1}{\omega_\pi(\vec{k}^2) [\omega_\pi(\vec{k}^2) + \omega_N(\vec{k}^2) - M_0]} \quad (22)$$

- On the domain for which the regularised integral has significant support, assume that M_0 is very much greater than all other mass scales.


$$\omega_N(\vec{k}^2) - M \approx \frac{\vec{k}^2}{2M} \quad (23)$$

- Then

$$\delta_F M_+^+ \approx -3g^2 \int \frac{d^3 k}{(2\pi)^3} \frac{\vec{k}^2}{8M^2 \omega_{\lambda_i}^2(\vec{k}^2)} \quad (24)$$

- So that

$$\frac{d^2 \delta_F M_+^+}{(dm_\pi^2)^2} \approx -\frac{3g^2}{4M^2} \int \frac{d^3 k}{(2\pi)^3} \frac{\vec{k}^2}{\omega_\pi^6(\vec{k}^2)} = -\frac{9}{128\pi} \frac{g^2}{M^2} \frac{1}{m_\pi}. \quad (25)$$



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

$$\delta_F M_+^+ = -3g^2 \int \frac{d^3 k}{(2\pi)^3} \frac{\omega_N(\vec{k}^2) - M_0}{4 \omega_N(\vec{k}^2)} \frac{1}{\omega_\pi(\vec{k}^2) [\omega_\pi(\vec{k}^2) + \omega_N(\vec{k}^2) - M_0]} \quad (26)$$

- On the domain for which the regularised integral has significant support, assume that M_0 is very much greater than all other mass scales.

$$\omega_N(\vec{k}^2) - M \approx \frac{\vec{k}^2}{2M} \quad (27)$$

- Then
$$\delta_F M_+^+ \approx -3g^2 \int \frac{d^3 k}{(2\pi)^3} \frac{\vec{k}^2}{8M^2 \omega_{\lambda_i}^2(\vec{k}^2)} \quad (28)$$

- So that
$$\frac{d^2 \delta_F M_+^+}{(dm_\pi^2)^2} \approx -\frac{3g^2}{4M^2} \int \frac{d^3 k}{(2\pi)^3} \frac{\vec{k}^2}{\omega_\pi^6(\vec{k}^2)} = -\frac{9}{128\pi} \frac{g^2}{M^2} \frac{1}{m_\pi}. \quad (29)$$

- Namely $\delta_F M_+^+ = -\frac{3}{32\pi} \frac{g^2}{M^2} m_\pi^3 + f_{(1)}^+(\lambda_1, \lambda_2) m_\pi^2 + f_{(0)}^+(\lambda_1, \lambda_2)$ where the last two terms express the necessary contribution from the regulator.



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)

Nucleon Self Energy: Chiral Limit

Hecht, *et al.*, nu-th/0201084

● Nucleon's self energy

$$\delta_F M_+^+ = -\frac{3}{32\pi} \frac{g^2}{M^2} m_\pi^3 + f_{(1)}^+(\lambda_1, \lambda_2) m_\pi^2 + f_{(0)}^+(\lambda_1, \lambda_2)$$



Nucleon Self Energy: Chiral Limit

Hecht, *et al.*, nu-th/0201084

- Nucleon's self energy

$$\delta_F M_+^+ = -\frac{3}{32\pi} \frac{g^2}{M^2} m_\pi^3 + f_{(1)}^+(\lambda_1, \lambda_2) m_\pi^2 + f_{(0)}^+(\lambda_1, \lambda_2)$$

- Given that $m_\pi^2 \propto \hat{m}$ in the neighbourhood of the chiral limit, the m_π^3 is nonanalytic in the current-quark mass on this domain.



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

- Nucleon's self energy

$$\delta_F M_+^+ = -\frac{3}{32\pi} \frac{g^2}{M^2} m_\pi^3 + f_{(1)}^+(\lambda_1, \lambda_2) m_\pi^2 + f_{(0)}^+(\lambda_1, \lambda_2)$$

- Given that $m_\pi^2 \propto \hat{m}$ in the neighbourhood of the chiral limit, the m_π^3 is nonanalytic in the current-quark mass on this domain.
- This is the **Leading Nonanalytic Contribution** much touted in effective field theory.
- Its form is completely fixed by chiral symmetry and the pattern of its dynamical breaking.

NB. Contribution from negative energy nucleon is $\propto \frac{1}{M^3}$.



Nucleon Self Energy: Chiral Limit

Hecht, et al., nu-th/0201084

- Nucleon's self energy

$$\delta_F M_+^+ = -\frac{3}{32\pi} \frac{g^2}{M^2} m_\pi^3 + f_{(1)}^+(\lambda_1, \lambda_2) m_\pi^2 + f_{(0)}^+(\lambda_1, \lambda_2)$$

- Given that $m_\pi^2 \propto \hat{m}$ in the neighbourhood of the chiral limit, the m_π^3 is nonanalytic in the current-quark mass on this domain.
 - This is the **Leading Nonanalytic Contribution** much touted in effective field theory.
 - Its form is completely fixed by chiral symmetry and the pattern of its dynamical breaking.

NB. Contribution from negative energy nucleon is $\propto \frac{1}{M^3}$.

- The **remaining** terms are regular in the current-quark mass. Their exact nature depends on the explicit form of regularisation procedure.



Nucleon Self Energy: Chiral Limit

Hecht, *et al.*, nu-th/0201084

- Nucleon's self energy

$$\delta_F M_+^+ = -\frac{3}{32\pi} \frac{g^2}{M^2} m_\pi^3 + f_{(1)}^+(\lambda_1, \lambda_2) m_\pi^2 + f_{(0)}^+(\lambda_1, \lambda_2)$$

- Given that $m_\pi^2 \propto \hat{m}$ in the neighbourhood of the chiral limit, the m_π^3 is nonanalytic in the current-quark mass on this domain.
- The **Leading Nonanalytic Contribution** is a model-independent result.



Nucleon Self Energy: Chiral Limit

Hecht, *et al.*, nu-th/0201084

- Nucleon's self energy

$$\delta_F M_+^+ = -\frac{3}{32\pi} \frac{g^2}{M^2} m_\pi^3 + f_{(1)}^+(\lambda_1, \lambda_2) m_\pi^2 + f_{(0)}^+(\lambda_1, \lambda_2)$$

- Given that $m_\pi^2 \propto \hat{m}$ in the neighbourhood of the chiral limit, the m_π^3 is nonanalytic in the current-quark mass on this domain.
- The **Leading Nonanalytic Contribution** is a model-independent result.
- **Unfortunately**, it is of limited relevance. In a calculation of the nucleon's mass, the actual value of the pion loop's contribution is almost completely determined by the regularisation dependent terms.
- It is essential for a framework to veraciously express the leading nonanalytic contribution . . . it serves as a check that DCSB is truly described.
- However, beyond that, one must accept that the world is complex.
 - The pion has a finite size. So does the nucleon.
 - These sizes set the mass-scale which determines the nucleon's mass shift.



Model pion-nucleon coupling

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Model pion-nucleon coupling

$$g_{PV}(P, k) = \frac{g}{2M} \exp(-(P - k)^2 / \Lambda^2)$$



Model pion-nucleon coupling

$$g_{PV}(P, k) = \frac{g}{2M} \exp(-(P - k)^2 / \Lambda^2)$$

• \mathcal{B} -Kernel

$$\int d\Omega_k g_{PV}^2((P - k)^2) \left[1 - \frac{2m_\pi^2}{(P - k)^2 + m_\pi^2} \right]$$

Clearly the sum of two independent terms.



Model pion-nucleon coupling

$$g_{PV}(P, k) = \frac{g}{2M} \exp(-(P - k)^2 / \Lambda^2)$$

- \mathcal{B} -Kernel

$$\int d\Omega_k g_{PV}^2((P - k)^2) \left[1 - \frac{2m_\pi^2}{(P - k)^2 + m_\pi^2} \right]$$

- First term can be evaluated exactly

$$\begin{aligned} \bar{g}_{PV}^2(P^2, k^2) &= \int d\Omega_k g_{PV}^2((P - k)^2) \\ &= \frac{g^2}{4M^2} e^{-2(P^2 + k^2)/\Lambda^2} \frac{\Lambda^2}{2Pk} I_1(4Pk/\Lambda^2), \end{aligned}$$



Model pion-nucleon coupling

$$g_{PV}(P, k) = \frac{g}{2M} \exp(-(P - k)^2 / \Lambda^2)$$

- \mathcal{B} -Kernel

$$\int d\Omega_k g_{PV}^2((P - k)^2) \left[1 - \frac{2m_\pi^2}{(P - k)^2 + m_\pi^2} \right]$$

- Second term can be approximated

$$\begin{aligned} \omega_{g^2}(P^2, k^2) &= 2m_\pi^2 \int d\Omega_k \frac{g_{PV}^2((P - k)^2)}{(P - k)^2 + m_\pi^2} \\ &\approx g_{PV}^2(|P - k|^2) \frac{2m_\pi^2}{a + \sqrt{a^2 - b^2}} \end{aligned}$$

- Reliable when analytic

structure of g_{PV} is not key to that of solution



Model pion-nucleon coupling

$$g_{PV}(P, k) = \frac{g}{2M} \exp(-(P - k)^2 / \Lambda^2)$$

- \mathcal{B} -Kernel

$$\int d\Omega_k g_{PV}^2((P - k)^2) \left[1 - \frac{2m_\pi^2}{(P - k)^2 + m_\pi^2} \right]$$

- Total Kernel:

$$\begin{aligned} &\approx \bar{g}_{PV}^2(P^2, k^2) - g_{PV}^2(|P^2 - k^2|) \frac{2m_\pi^2}{a + \sqrt{a^2 - b^2}}, \\ &=: \bar{g}_{PV}^2(P^2, k^2) - \tilde{g}_{PV}^2(P^2, k^2) \frac{2m_\pi^2}{a + \sqrt{a^2 - b^2}}, \end{aligned}$$

- Analytic structure is transparent



Nucleon's self energy and mass shift



Nucleon's self energy and mass shift

- Solve DSE Nonperturbatively

$$M_D^2 \mathcal{A}^2(-M_D^2) = [M + \mathcal{B}(-M_D^2)]^2$$

$$\delta M = M_D - M$$



Nucleon's self energy and mass shift

- Solve DSE Nonperturbatively

$$M_D^2 \mathcal{A}^2(-M_D^2) = [M + \mathcal{B}(-M_D^2)]^2$$
$$\delta M = M_D - M$$

- Vector self energy



Nucleon's self energy and mass shift

- Solve DSE Nonperturbatively

$$M_D^2 \mathcal{A}^2(-M_D^2) = [M + \mathcal{B}(-M_D^2)]^2$$
$$\delta M = M_D - M$$

- Scalar self energy




Nucleon's self energy and mass shift

- Solve DSE Nonperturbatively

$$M_D^2 \mathcal{A}^2(-M_D^2) = [M + \mathcal{B}(-M_D^2)]^2$$

$$\delta M = M_D - M$$

	(Λ, Λ_N)	(Λ, Λ_N)	(Λ, Λ_N)
	$(0.9, \infty)$	$(0.9, 1.5)$	$(0.9, 2.0)$
$-\delta M$ (MeV)	222	61	99



Nucleon's self energy and mass shift

- Solve DSE Nonperturbatively

$$M_D^2 \mathcal{A}^2(-M_D^2) = [M + \mathcal{B}(-M_D^2)]^2$$

$$\delta M = M_D - M$$

	(Λ, Λ_N)	(Λ, Λ_N)	(Λ, Λ_N)
	$(0.9, \infty)$	$(0.9, 1.5)$	$(0.9, 2.0)$
$-\delta M$ (MeV)	222	61	99

- No suppression for nucleon off-shell in self-energy loop;
i.e, $g_{PV}((P - k^2), P^2, k^2)$

Neglected this dependence



Nucleon's self energy and mass shift

- Solve DSE Nonperturbatively

$$M_D^2 \mathcal{A}^2(-M_D^2) = [M + \mathcal{B}(-M_D^2)]^2$$

$$\delta M = M_D - M$$

	(Λ, Λ_N)	(Λ, Λ_N)	(Λ, Λ_N)
	$(0.9, \infty)$	$(0.9, 1.5)$	$(0.9, 2.0)$
$-\delta M$ (MeV)	222	61	99



$$g_{PV}(P^2, k^2, P \cdot k) = \frac{g}{2M} e^{-(P-k)^2/\Lambda^2} e^{-(P^2+M^2+k^2+M^2)/\Lambda_N^2}$$

- Correct on-shell limit:

$$g_{PV}(P^2 = -M^2, k^2 = -M^2, (P - k)^2 = 0) = \frac{g}{2M}$$

Nucleon's self energy and mass shift

- Solve DSE Nonperturbatively

$$M_D^2 \mathcal{A}^2(-M_D^2) = [M + \mathcal{B}(-M_D^2)]^2$$

$$\delta M = M_D - M$$

Range from meson
exchange model phen.

	(Λ, Λ_N)	(Λ, Λ_N)	(Λ, Λ_N)
	$(0.9, \infty)$	$(0.9, 1.5)$	$(0.9, 2.0)$
$-\delta M$ (MeV)	222	61	99



$$g_{PV}(P^2, k^2, P \cdot k) = \frac{g}{2M} e^{-(P-k)^2/\Lambda^2} e^{-(P^2+M^2+k^2+M^2)/\Lambda_N^2}$$

$\Lambda_N \rightarrow \infty \Rightarrow$ pointlike nucleon

Pion loop's effect

- Nonpointlike πN -loop
 - ... reduces nucleon's mass by ~ 100 MeV



Pion loop's effect

- Nonpointlike πN -loop
 - ... reduces nucleon's mass by ~ 100 MeV
- There's also a $\pi \Delta$ -loop
 - ... reduces nucleon's mass by not more than 100 MeV



Pion loop's effect

- Nonpointlike πN -loop
 - ... reduces nucleon's mass by ~ 100 MeV
- There's also a $\pi \Delta$ -loop
 - ... reduces nucleon's mass by not more than 100 MeV

$$-\delta M_N \sim 200 \text{ MeV}$$



Pion loop's effect

- Nonpointlike πN -loop
 - ... reduces nucleon's mass by ~ 100 MeV
- There's also a $\pi \Delta$ -loop
 - ... reduces nucleon's mass by not more than 100 MeV
- $-\delta M_N \sim 200$ MeV
- Qualitative effect of this?



Too much of a good thing

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Too much of a good thing

- Refit Faddeev model parameters,
allowing for heavier “quark-core” mass



Too much of a good thing

	ω_{0+}	ω_{1+}	M_N	M_Δ	ω_{f_1}	ω_{f_2}	R
0^+	0.45	-	1.44	-	0.36	0.35	2.32
$0^+ \text{ \& } 1^+$	0.45	1.36	1.14	1.33	0.44	0.36	0.54
0^+	0.64	-	1.59	-	0.39	0.41	1.28
$0^+ \text{ \& } 1^+$	0.64	1.19	0.94	1.23	0.49	0.44	0.25



50% reduction in role of axial-vector diquark

Too much of a good thing

	ω_{0+}	ω_{1+}	M_N	M_Δ	ω_{f_1}	ω_{f_2}	R
0^+	0.45	-	1.44	-	0.36	0.35	2.32
$0^+ \text{ \& } 1^+$	0.45	1.36	1.14	1.33	0.44	0.36	0.54
0^+	0.64	-	1.59	-	0.39	0.41	1.28
$0^+ \text{ \& } 1^+$	0.64	1.19	0.94	1.23	0.49	0.44	0.25



- 50% reduction in role of axial-vector diquark
- 10% increase in role of scalar diquark

Too much of a good thing

	ω_{0+}	ω_{1+}	M_N	M_Δ	ω_{f_1}	ω_{f_2}	R
0^+	0.45	-	1.44	-	0.36	0.35	2.32
$0^+ \text{ \& } 1^+$	0.45	1.36	1.14	1.33	0.44	0.36	0.54
0^+	0.64	-	1.59	-	0.39	0.41	1.28
$0^+ \text{ \& } 1^+$	0.64	1.19	0.94	1.23	0.49	0.44	0.25

Unsurprisingly:

Requiring **Exact Fit** to N , Δ masses
 with **only** q , $(qq)_{JP}$ Degrees of Freedom
 \Rightarrow **Forces** 1^+ to mimic, **in part**, effect of π



Pseudoscalar mesons and Form Factors

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Pseudoscalar mesons and Form Factors

- Light mass of pseudoscalar mesons means they play a very important role in many aspects of hadron physics.



Pseudoscalar mesons and Form Factors

- Light mass of pseudoscalar mesons means they play a very important role in many aspects of hadron physics.
- Indeed, no approach to low-energy hadron physics that does not explicitly account for pseudoscalar meson degrees of freedom can be valid.



Pseudoscalar mesons and Form Factors

- Light mass of pseudoscalar mesons means they play a very important role in many aspects of hadron physics.
- Indeed, no approach to low-energy hadron physics that does not explicitly account for pseudoscalar meson degrees of freedom can be valid.
- Another example . . . pseudoscalar mesons also contribute materially to form factors.



Pseudoscalar mesons and Form Factors

- Light mass of pseudoscalar mesons means they play a very important role in many aspects of hadron physics.
- Indeed, no approach to low-energy hadron physics that does not explicitly account for pseudoscalar meson degrees of freedom can be valid.
- Another example . . . pseudoscalar mesons also contribute materially to form factors.
- Illustrate with $\gamma N \rightarrow \Delta$ transition form factor. Focus on the M1 (spin-flip) form factor, $G_M^*(Q^2)$.



Harry Lee

Pions and Form Factors



[First](#)

[Contents](#)

[Back](#)

[Conclusion](#)

Pions and Form Factors


- Dynamical coupled-channels model . . . Analyzed extensive JLab data . . . Completed a study of the $\Delta(1236)$
 - *Meson Exchange Model for πN Scattering and $\gamma N \rightarrow \pi N$ Reaction*, T. Sato and T.-S. H. Lee, Phys. Rev. C **54**, 2660 (1996)
 - *Dynamical Study of the Δ Excitation in $N(e, e'\pi)$ Reactions*, T. Sato and T.-S. H. Lee, Phys. Rev. C **63**, 055201/1-13 (2001)



Pions and Form Factors

- Dynamical coupled-channels model ... Analyzed extensive JLab data ... Completed a study of the $\Delta(1236)$
 - *Meson Exchange Model for πN Scattering and $\gamma N \rightarrow \pi N$ Reaction*, T. Sato and T.-S. H. Lee, Phys. Rev. C **54**, 2660 (1996)
 - *Dynamical Study of the Δ Excitation in $N(e, e'\pi)$ Reactions*, T. Sato and T.-S. H. Lee, Phys. Rev. C **63**, 055201/1-13 (2001)
- Pion cloud effects are large in the low Q^2 region.

Ratio of the M1 form factor in $\gamma N \rightarrow \Delta$ transition and proton dipole form factor G_D . Solid curve is $G_M^(Q^2)/G_D(Q^2)$ including pions; Dotted curve is $G_M(Q^2)/G_D(Q^2)$ without pions.*




Pions and Form Factors

- Dynamical coupled-channels model ... Analyzed extensive JLab data ... Completed a study of the $\Delta(1236)$
 - *Meson Exchange Model for πN Scattering and $\gamma N \rightarrow \pi N$ Reaction*, T. Sato and T.-S. H. Lee, Phys. Rev. C **54**, 2660 (1996)
 - *Dynamical Study of the Δ Excitation in $N(e, e'\pi)$ Reactions*, T. Sato and T.-S. H. Lee, Phys. Rev. C **63**, 055201/1-13 (2001)
- Pion cloud effects are large in the low Q^2 region.

Ratio of the M1 form factor in $\gamma N \rightarrow \Delta$ transition and proton dipole form factor G_D . Solid curve is $G_M^*(Q^2)/G_D(Q^2)$ including pions; Dotted curve is $G_M(Q^2)/G_D(Q^2)$ without pions.

Quark Core

- Responsible for only 2/3 of result at small Q^2
- Dominant for $Q^2 > 2 - 3 \text{ GeV}^2$



Results: Nucleon and Δ Masses



Results: Nucleon and Δ Masses

Mass-scale parameters (in GeV)
for the scalar and axial-vector
diquark correlations, fixed by
fitting nucleon and Δ masses



Set A – fit to the actual masses was required; whereas for
Set B – fitted mass was offset to allow for “ π -cloud” contributions



Argonne
NATIONAL
LABORATORY

set	M_N	M_Δ	m_{0+}	m_{1+}	ω_{0+}	ω_{1+}
A	0.94	1.23	0.63	0.84	$0.44=1/(0.45 \text{ fm})$	$0.59=1/(0.33 \text{ fm})$
B	1.18	1.33	0.80	0.89	$0.56=1/(0.35 \text{ fm})$	$0.63=1/(0.31 \text{ fm})$

● $m_{1+} \rightarrow \infty$: $M_N^A = 1.15 \text{ GeV}$; $M_N^B = 1.46 \text{ GeV}$

Results: Nucleon and Δ Masses

Mass-scale parameters (in GeV) for the scalar and axial-vector diquark correlations, fixed by fitting nucleon and Δ masses



Set A – fit to the actual masses was required; whereas for **Set B** – fitted mass was offset to allow for “ π -cloud” contributions



Argonne
NATIONAL
LABORATORY

set	M_N	M_Δ	m_{0+}	m_{1+}	ω_{0+}	ω_{1+}
A	0.94	1.23	0.63	0.84	$0.44=1/(0.45 \text{ fm})$	$0.59=1/(0.33 \text{ fm})$
B	1.18	1.33	0.80	0.89	$0.56=1/(0.35 \text{ fm})$	$0.63=1/(0.31 \text{ fm})$


● $m_{1+} \rightarrow \infty$: $M_N^A = 1.15 \text{ GeV}$; $M_N^B = 1.46 \text{ GeV}$

● Axial-vector diquark provides significant attraction

Results: Nucleon and Δ Masses

Mass-scale parameters (in GeV) for the scalar and axial-vector diquark correlations, fixed by fitting nucleon and Δ masses

Set A – fit to the actual masses was required; whereas for **Set B** – fitted mass was offset to allow for “ π -cloud” contributions



Argonne
NATIONAL
LABORATORY

set	M_N	M_Δ	m_{0+}	m_{1+}	ω_{0+}	ω_{1+}
A	0.94	1.23	0.63	0.84	$0.44=1/(0.45 \text{ fm})$	$0.59=1/(0.33 \text{ fm})$
B	1.18	1.33	0.80	0.89	$0.56=1/(0.35 \text{ fm})$	$0.63=1/(0.31 \text{ fm})$

● $m_{1+} \rightarrow \infty$: $M_N^A = 1.15 \text{ GeV}$; $M_N^B = 1.46 \text{ GeV}$

● **Constructive Interference:** 1^{++} -diquark + $\partial_\mu \pi$

Craig Roberts: Dyson Schwinger Equations and QCD

25th Students' Workshop on Electromagnetic Interactions, 31/08 – 05/09, 2008. ... – p. 30/38

First

Contents

Back

Conclusion

Nucleon-Photon Vertex

[First](#)[Contents](#)[Back](#)[Conclusion](#)

M. Oettel, M. Pichowsky
and L. von Smekal, nu-th/9909082

6 terms . . .

Nucleon-Photon Vertex








constructed systematically . . . current conserved automatically
for on-shell nucleons described by Faddeev Amplitude




6 terms ...

Nucleon-Photon Vertex

constructed systematically ... current conserved automatically
for on-shell nucleons described by Faddeev Amplitude




Form Factor Ratio: *GE/GM*



Form Factor Ratio:

GE/GM

- Combine these elements ...




Form Factor Ratio:

GE/GM

● Combine these elements ...


● Dressed-Quark Core



● Combine these elements ...


● Dressed-Quark Core

● *Ward-Takahashi*
Identity preserving
current




● Combine these elements ...

- Dressed-Quark Core
- *Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution



● Combine these elements ...


- Dressed-Quark Core
- *Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution



● Combine these elements ...


- Dressed-Quark Core
- *Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution

● All parameters fixed in
other applications ... **Not** varied.



- Combine these elements ...

- Dressed-Quark Core
- Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution




- All parameters fixed in
other applications ... **Not** varied.

- Agreement with Pol. Trans. data at $Q^2 \gtrsim 2 \text{ GeV}^2$



- Combine these elements ...

- Dressed-Quark Core
- Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution




- All parameters fixed in
other applications ... **Not** varied.

- Agreement with Pol. Trans. data at $Q^2 \gtrsim 2 \text{ GeV}^2$
- Correlations in Faddeev amplitude – quark orbital
angular momentum – essential to that agreement



- Combine these elements ...

- Dressed-Quark Core
- Ward-Takahashi*
Identity preserving
current
- Anticipate and
Estimate Pion
Cloud's Contribution



- All parameters fixed in
other applications ... **Not** varied.

- Agreement with Pol. Trans. data at $Q^2 \gtrsim 2 \text{ GeV}^2$
- Correlations in Faddeev amplitude – quark orbital
angular momentum – essential to that agreement
- Predict Zero at $Q^2 \approx 6.5 \text{ GeV}^2$



Density profile of charge and magnetisation

[First](#)[Contents](#)[Back](#)[Conclusion](#)


Density profile of charge and magnetisation

- Proton's Electromagnetic Form Factor
 - Appearance of a zero in $G_E(Q^2)$ – Completely Unexpected




Density profile of charge and magnetisation

- Proton's Electromagnetic Form Factor
 - Appearance of a zero in $G_E(Q^2)$ – Completely Unexpected



Density profile of charge and magnetisation

- Proton's Electromagnetic Form Factor
 - Appearance of a zero in $G_E(Q^2)$ – Completely Unexpected



● However, Current Density remains peaked at $r = 0$!

Craig Roberts: Dyson Schwinger Equations and QCD

25th Students' Workshop on Electromagnetic Interactions, 31/08 – 05/09, 2008. ... – p. 33/38



Density profile of charge and magnetisation

- Proton's Electromagnetic Form Factor
 - Appearance of a zero in $G_E(Q^2)$ – Completely Unexpected
- Wave Function is complex and correlated mix of virtual particles and antiparticles: s –, p – and d –waves



Density profile of charge and magnetisation

- Proton's Electromagnetic Form Factor
 - Appearance of a zero in $G_E(Q^2)$ – Completely Unexpected
- Wave Function is complex and correlated mix of virtual particles and antiparticles: s –, p – and d –waves
- *Simple independent-particle three-quark bag-model picture is profoundly incorrect*



Improved current

[First](#)[Contents](#)[Back](#)[Conclusion](#)

Improved current

- Composite axial-vector diquark correlation
 - Electromagnetic current can be complicated
 - Limited constraints on large- Q^2 behaviour



Improved current

- Composite axial-vector diquark correlation
 - Electromagnetic current can be complicated
 - Limited constraints on large- Q^2 behaviour
 - Improved performance of code
 - Implemented corrections so that large- Q^2 behaviour of form factors could be reliably calculated
 - Exposed two weaknesses in rudimentary *Ansatz*



Improved current

- Composite axial-vector diquark correlation
 - Improved performance of code
 - Implemented corrections so that large- Q^2 behaviour of form factors could be reliably calculated
 - Exposed two weaknesses in rudimentary *Ansatz*
 - Diquark effectively pointlike to hard probe
 - Didn't account for diquark being off-shell in recoil




Improved current

- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%



Improved current


- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%



Improved current

- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%


● Proton – zero shifted:
 $6.5 \rightarrow 8.0 \text{ GeV}^2$



Improved current

- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%

- **Proton** – zero shifted:
 $6.5 \rightarrow 8.0 \text{ GeV}^2$



Improved current

- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%


● **Proton** – zero shifted:

$$6.5 \rightarrow 8.0 \text{ GeV}^2$$

Neutron – peak shifted:

$$7.5 \rightarrow 5.0 \text{ GeV}^2$$

& now predict zero
a little above 11 GeV^2



Improved current

- Composite axial-vector diquark correlation
- Minor but material improvements to current
 - Introduce form factor: radius 0.8 fm
 - Increase recoil mass by 10%


● **Proton** – zero shifted:

$6.5 \rightarrow 8.0 \text{ GeV}^2$

Neutron – peak shifted:

$7.5 \rightarrow 5.0 \text{ GeV}^2$

& now predict zero
a little above 11 GeV^2






Pion Cloud

F_1 – neutron


- Comparison between Faddeev equation result and Kelly's parametrisation



Pion Cloud

F_1 – neutron


- Comparison between Faddeev equation result and Kelly's parametrisation
- Faddeev equation set-up to describe dressed-quark core



Pion Cloud

F_1 – neutron

- Comparison between Faddeev equation result and Kelly's parametrisation
- Faddeev equation set-up to describe dressed-quark core
- Pseudoscalar meson cloud (and related effects) significant for $Q^2 \lesssim 3 - 4 M_N^2$





Epilogue

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Epilogue

[First](#)[Contents](#)[Back](#)[Conclusion](#)



Epilogue

- DCSB exists in QCD.





Epilogue

- DCSB exists in QCD.
- It is manifest in dressed propagators and vertices





Epilogue

- DCSB exists in QCD.
 - It is manifest in dressed propagators and vertices
 - It predicts, amongst other things, that
 - light current-quarks become heavy constituent-quarks
 - pseudoscalar mesons are unnaturally light
 - pseudoscalar mesons couple unnaturally strongly to light-quarks
 - pseudoscalar mesons couple unnaturally strongly to the lightest baryons





Epilogue

- DCSB exists in QCD.
 - It is manifest in dressed propagators and vertices
 - It predicts, amongst other things, that
 - light current-quarks become heavy constituent-quarks
 - pseudoscalar mesons are unnaturally light
 - pseudoscalar mesons couple unnaturally strongly to light-quarks
 - pseudoscalar mesons couple unnaturally strongly to the lightest baryons
 - It impacts dramatically upon observables.





Epilogue

- Form Factors - progress anticipated in near- to medium-term
 - Quantifying pseudoscalar meson “cloud” effects





Epilogue

- Form Factors - progress anticipated in near- to medium-term
 - Quantifying pseudoscalar meson “cloud” effects
 - Locating and explaining the transition from nonp-QCD to p-QCD in the pion and nucleon electromagnetic form factors





Epilogue

- Form Factors - progress anticipated in near- to medium-term
 - Quantifying pseudoscalar meson “cloud” effects
 - Locating and explaining the transition from nonp-QCD to p-QCD in the pion and nucleon electromagnetic form factors
 - Explaining the high Q^2 behavior of the proton form factor ratio in the space-like region





Epilogue

- Form Factors - progress anticipated in near- to medium-term
 - Quantifying pseudoscalar meson “cloud” effects
 - Locating and explaining the transition from nonp-QCD to p-QCD in the pion and nucleon electromagnetic form factors
 - Explaining the high Q^2 behavior of the proton form factor ratio in the space-like region
 - Detailing broadly the role of two-photon exchange contributions






nothing!

Epilogue

- Form Factors - progress anticipated in near- to medium-term
 - Quantifying pseudoscalar meson “cloud” effects
 - Locating and explaining the transition from nonp-QCD to p-QCD in the pion and nucleon electromagnetic form factors
 - Explaining the high Q^2 behavior of the proton form factor ratio in the space-like region
 - Detailing broadly the role of two-photon exchange contributions
 - Explaining relationship between parton properties on the light-front and rest frame structure of hadrons



Vector piece of nucleon's self energy




$$M = 0.94, m_\pi = 0.14$$

$$\Lambda = 0.9 \text{ GeV}, g_A = 1$$



Vector piece of nucleon's self energy



- Solid line: One-loop; numerically evaluated, exact (numerical) kernel




$$M = 0.94, m_\pi = 0.14$$


$$\Lambda = 0.9 \text{ GeV}, g_A = 1$$



Vector piece of nucleon's self energy



- Solid line: One-loop; numerically evaluated, exact (numerical) kernel
- Dotted line: One-loop; algebraic; code test




$$M = 0.94, m_\pi = 0.14$$

$$\Lambda = 0.9 \text{ GeV}, g_A = 1$$



Vector piece of nucleon's self energy




- Solid line: One-loop; numerically evaluated, exact (numerical) kernel
- Dotted line: One-loop; algebraic; code test
- Dashed line: Fully self-consistent numerical solution; extra pion loops do little




$$M = 0.94, m_\pi = 0.14$$
$$\Lambda = 0.9 \text{ GeV}, g_A = 1$$



Vector piece of nucleon's self energy




- Solid line: One-loop; numerically evaluated, exact (numerical) kernel
- Dotted line: One-loop; algebraic; code test
- Dot-dashed line: Fully self-consistent numerical solution; continuation to timelike region



$$M = 0.94, m_\pi = 0.14$$
$$\Lambda = 0.9 \text{ GeV}, g_A = 1$$



Scalar piece of nucleon's self energy




$$M = 0.94, m_\pi = 0.14$$

$$\Lambda = 0.9 \text{ GeV}, g_A = 1$$



Scalar piece of nucleon's self energy




- Solid line: One-loop; numerically evaluated, exact (numerical) kernel



$$M = 0.94, m_\pi = 0.14$$
$$\Lambda = 0.9 \text{ GeV}, g_A = 1$$



Scalar piece of nucleon's self energy



- Solid line: One-loop; numerically evaluated, exact (numerical) kernel
- Dotted line: One-loop; algebraic; code test




$$M = 0.94, m_\pi = 0.14$$

$$\Lambda = 0.9 \text{ GeV}, g_A = 1$$



Scalar piece of nucleon's self energy




- Solid line: One-loop; numerically evaluated, exact (numerical) kernel
- Dotted line: One-loop; algebraic; code test
- Dashed line: Fully self-consistent numerical solution; extra pion loops do little



$$M = 0.94, m_\pi = 0.14$$
$$\Lambda = 0.9 \text{ GeV}, g_A = 1$$



Scalar piece of nucleon's self energy



- Solid line: One-loop; numerically evaluated, exact (numerical) kernel
- Dotted line: One-loop; algebraic; code test
- Dot-dashed line: Fully self-consistent numerical solution; continuation to timelike region



$$M = 0.94, m_\pi = 0.14$$
$$\Lambda = 0.9 \text{ GeV}, g_A = 1$$

